arXiv:1505.00747v1 [math.DG] 4 May 2015

THE ASYMPTOTIC BEHAVIOR OF THE MONODROMY REPRESENTATIONS OF THE ASSOCIATED FAMILIES OF COMPACT CMC SURFACES

SEBASTIAN HELLER

Abstract. Constant mean curvature (CMC) surfaces in space forms can be described by their associated $\mathbb{C}^*$-family of flat $\mathrm{SL}(2,\mathbb{C})$-connections ∇^λ. In this paper we consider the asymptotic behavior (for $\lambda \to 0$) of the gauge equivalence classes of ∇^λ for compact CMC surfaces of genus $g \geq 2$. We prove (under the assumption of simple umbilics) that the asymptotic behavior of the traces of the monodromy representation of ∇^λ determines the conformal type as well as the Hopf differential locally in the Teichmüller space.

1. Introduction

CMC surfaces in space forms are given by their associated $\mathbb{C}^*$-family of flat $\mathrm{SL}(2,\mathbb{C})$-connections ∇^λ. For simply connected surfaces, all flat connections are trivial, and the associated family of gauge equivalence classes $[\nabla^\lambda]$ does not contain any information. This is in stark contrast to compact surfaces of genus $g \geq 1$. For CMC tori, it was shown by Hitchin [9] that the conformal type of the torus (as well as its energy) are determined by the asymptotic behavior of $[\nabla^\lambda]$ as $\lambda \to 0$. Similarly, it was shown in [8] that the energy of Lawson symmetric CMC surfaces of genus 2 can also be recovered from the asymptotic behavior of $[\nabla^\lambda]$. In this paper, we prove that also the conformal type of a compact CMC surface of genus $g \geq 2$ with simple umbilics is determined by the asymptotic behavior of the associated family of gauge equivalence classes $[\nabla^\lambda]$ locally in the Teichmüller space: The trace of the monodromy of ∇^λ along curves γ (satisfying a certain condition) give the period of the square root of the Hopf differential along γ, see formula (2.5). Then, we reprove that the periods of the square root of a holomorphic quadratic differential with simple zeros (defined on the so-called Hitchin covering) determine the Riemann surface structure locally in the Teichmüller space. We then apply this observation to prove our main theorem.

2. The family of flat connections and its asymptotic monodromy

Let $f\colon M \to S^3$ be a conformal immersion of constant mean curvature H from a compact Riemann surface to the round S^3. The immersion f is determined by its associated $\mathbb{C}^*$-family ∇^λ of flat $\mathrm{SL}(2,\mathbb{C})$ connections on the trivial $\mathbb{C}^2$ bundle over M, see [9, 1]:

Theorem 1 ([9, 1]). *For a conformal CMC immersion $f\colon M \to S^3$ there exists an associated family of flat $\mathrm{SL}(2,\mathbb{C})$-connections*

$$\lambda \in \mathbb{C}^* \mapsto \nabla^\lambda = \nabla + \lambda^{-1}\Phi - \lambda\Phi^* \tag{2.1}$$

Date: May 5, 2015.

The author would like to thank Aaron Gerding for many interesting discussions. Author supported by DFG HE 6829/1-2.

on a complex bundle $V \to M$ *of rank* 2, *where* Φ *is a nilpotent and nowhere vanishing complex linear* 1*-form and* Φ^* *is its adjoint with respect to a unitary metric. The connections* ∇^λ *are unitary for* $S^1 \subset \mathbb{C}^*$ *and trivial for* $\lambda_1 \neq \lambda_2 \in S^1$.

Conversely, for every $\mathbb{C}^*$*-family of flat connections satisfying the properties listed above, the* $SU(2)$*-valued gauge transformation between the trivial connections* ∇^{λ_1} *and* ∇^{λ_2} *is a CMC immersion into* $S^3 = SU(2)$ *with mean curvature* $H = i\frac{\lambda_1+\lambda_2}{\lambda_1-\lambda_2}$.

Because of the following theorem we cannot apply the methods developed by Hitchin [9] in order to study CMC surfaces of higher genus.

Theorem 2 ([7]). *For compact CMC surfaces of genus* $g \geq 2$ *the flat* $SL(2,\mathbb{C})$*-connections* ∇^λ *are irreducible for generic spectral parameter* $\lambda \in \mathbb{C}^*$.

2.1. **Monodromy.** Because Φ in Theorem 1 is nilpotent and nowhere vanishing, there exists a holomorphic line bundle $S^* = \ker \Phi \subset V$ which automatically satisfies $(S^*)^2 = K^*$, i.e., S^* is the dual of a holomorphic spin bundle S. With respect to the unitary decomposition $V = S^* \oplus S$ the connections ∇^λ are given by

$$\nabla^\lambda = \begin{pmatrix} \nabla^{spin^*} & -\frac{i}{2}Q^* \\ -\frac{i}{2}Q & \nabla^{spin} \end{pmatrix} + \lambda^{-1}\begin{pmatrix} 0 & \frac{1}{2} \\ 0 & 0 \end{pmatrix} - \lambda\begin{pmatrix} 0 & 0 \\ i\,\mathrm{vol} & 0 \end{pmatrix}, \tag{2.2}$$

see [7, 8]. In (2.2) Q is a holomorphic quadratic differential – the Hopf differential of the CMC surface – and ∇^S is the spin connection of the induced Levi-Civita connection. Let $\zeta^2 = \lambda$ and consider the ζ-dependent gauge transformations

$$g = \begin{pmatrix} \frac{1}{\sqrt{\zeta}} & 0 \\ 0 & \sqrt{\zeta} \end{pmatrix}$$

with respect to the decomposition $V = S^* \oplus S$. Then,

$$\nabla^\zeta := \nabla^\lambda.g = \tilde{\nabla} + \zeta^{-1}\begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{i}{2}Q & 0 \end{pmatrix} - \zeta\begin{pmatrix} 0 & -\frac{i}{2}Q^* \\ i\,\mathrm{vol} & 0 \end{pmatrix}, \tag{2.3}$$

is the twisted family of flat connections. For $\zeta^2 = \lambda$ the traces of the monodromies of ∇^ζ and ∇^λ along any closed curve are the same.

Instead of working with ∇^λ or ∇^ζ we will work with another family of flat connections: Consider the Hopf differential Q of the CMC surface of genus $g \geq 2$ and assume that it has only simple zeros. Then there exists a double covering $\hat{M} \to M$ (the Hitchin covering [10]) branched over the umbilics, i.e., the zeros of Q. $\hat{M}$ has genus $4g-3$ and there exists a holomorphic 1-form ω on $\hat{M}$ such that $\frac{i}{4}Q = -\omega^2$. Note that in [10] Q is pulled back as a section and not as a quadratic differential. Nevertheless, it follows easily that the pullback of Q as a quadratic differential has a globally well-defined square root which is a holomorphic 1-form on the Hitchin covering.

The holomorphic Higgs field $\Phi = \begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{i}{2}Q & 0 \end{pmatrix}$ (with respect to $\bar\partial^{\tilde{\nabla}}$) diagonalizes on $\hat{M}$ [10], i.e., there is a decomposition $V = L \oplus L^* \to \hat{M}$ such that

$$\Phi = \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix},$$

where Φ is the pull-back as an endomorphism-valued 1-form. In the following, we will work with the pull-back connections $\hat{\nabla}^\zeta$ of ∇^ζ on $V = L \oplus L^* \to \hat{M}$.

In order to obtain informations from the asymptotic behavior of the monodromy of $\hat{\nabla}^\zeta$ (respectively ∇^λ) we need to recall basic knowledge from the asymptotic analysis of linear ordinary differential equations. For a more extensive treatment of this topic we refer to [13, 2]. A short treatment appropriate for our purposes is given in Appendix C in [4]: Let $I = (\varphi_0; \varphi_1) \subset \mathbb{R}$ and consider the sector

$$S := \{te^{i\varphi} \mid t \in \mathbb{R}^{>0}, \varphi \in I\}.$$

Let $\gamma\colon [0;1] \to \hat{M}$ be a closed curve such that

$$\mathrm{Re}(e^{i\varphi}\omega(\gamma'(t))) > 0 \tag{2.4}$$

for all directions $e^{i\varphi} \in S^1$ with $\varphi \in I$ and all $t \in [0;1]$. Then the trace of the monodromy along γ satisfies

$$\mathrm{tr}(M(\hat{\nabla}^\zeta, \gamma)) \exp\left(-\tfrac{1}{\zeta}\int_\gamma \omega\right) \xrightarrow[\frac{1}{\zeta}\in S,\, \zeta\to 0]{} \mathit{const} \neq 0 \tag{2.5}$$

for some $\mathit{const} \in \mathbb{C} \setminus \{0\}$. Hence, we obtain:

Proposition 2.1. *For closed curves γ satisfying* (2.4) *the value of the period $\int_\gamma \omega$ is determined by the asymptotic behavior for $\lambda \to 0$ of the trace of the monodromy representation of ∇^λ.*

3. The Riemann surface structure

Let M_t, $t \in (-\delta, \delta)$, be a (smooth) family of compact Riemann surfaces of genus g, and let Q_t be a family of holomorphic quadratic differentials with respect to M_t. Assume that the differentials Q_t have only simple zeros, and consider the family $\pi_t\colon \hat{M}_t \to M_t$ of Hitchin coverings as well as the corresponding (holomorphic) involutions σ_t interchanging the sheets. We can identify the smooth surfaces $\hat{M} = \hat{M}_t$ in such a way that all σ_t are the same (smooth map $\sigma\colon \hat{M} \to \hat{M}$). In this situation we get:

Lemma 3.1. *Assume that on the Hitchin covering $\hat{M} \to M$ the periods of $\omega_t = \sqrt{Q}_t \in H^0(\hat{M}_t, K_{\hat{M}_t})$ are constant in t. Then the Riemann surfaces M_t are equivalent. Moreover, the Hopf differentials Q_t are also equivalent.*

Remark 3.1. It is a well-known fact (see [11] or §1.1. in [12]) that the space of Riemann surfaces equipped with holomorphic differentials ω having a prescribed number of zeros of prescribed order has local coordinates given by the so-called relative periods $\int_{\gamma_k} \omega$, where the curves γ_k, $k = 1, ..$, generate the first homology group relative to the set of zeros of ω. We briefly reprove this fact in the situation of this paper along with the necessary computation of the "half-periods" $\int_\gamma \omega$ for curves connecting zeros of ω.

Proof. Consider the smooth surface $\hat{M}$ together with the family of closed complex valued 1-forms ω_t. They are holomorphic with respect to a t-dependent Riemann surface structure J_t, and their zeros are all of order 2. Hence, we can assume that the zeros coincide on the smooth surface $\hat{M}$. Moreover, we can also assume without loss of generality that in a local neighborhood $U \subset \hat{M}$ of its common zeros, the 1-forms coincide $(\omega_t)_{|U} = (\omega_0)_{|U}$. As the periods of ω_t are independent of t we get

$$\omega_t = \omega_0 + df_t$$

for some complex-valued function $f_t\colon \hat{M} \to \mathbb{C}$ depending smoothly on t. Moreover, the differential $df_t \in \Omega^1(\hat{M}, \mathbb{C})$ vanishes in the neighborhood U of the zeros of ω. In order to apply the Moser trick to deform the forms ω_t to ω_0, we first need to establish that the values of $\frac{\partial f_t}{\partial t}$ near the zeros of ω_t are 0. Recall that we have already identified the involutions $\sigma_t = \sigma\colon \hat{M} \to \hat{M}$. Let γ be a curve connecting two zeros of ω_t and consider the closed curve $\Gamma = \sigma(\gamma^{-1}) \circ \gamma$. Because $\sigma^*\omega_t = -\omega_t$ for all t we obtain

$$2\int_\gamma \omega_t = \int_\Gamma \omega_t = \int_\Gamma \omega_0 = 2\int_\gamma \omega_0.$$

Therefore, the values of f_t at any zero of ω_t are the same, and without loss of generality we have that they are 0 independently of t. Next we consider the closed 1-form

$$\Omega = \omega_0 + df_t + \frac{\partial f_t}{\partial t}dt = \omega_t + \frac{\partial f_t}{\partial t}dt$$

on $\hat{M} \times (-\delta, \delta)$. We want to show that there exists a (real) vector field X on $\hat{M} \times (-\delta, \delta)$ such that $\Omega(X) = 0$ and $dt(X) = 1$. We already know that f_t is 0 in the neighborhood U of the zeros of ω_t for all $t \in (-\delta, \delta)$. Therefore, the existence and uniqueness of X follows from the fact that for all $p \in \hat{M} \setminus \{\text{zeros of } \omega_t\}$ the linear map $(\omega_t)_p\colon T_p\hat{M} \to \mathbb{C}$ is an isomorphism. Clearly, X is complete. Then

$$\mathcal{L}_X\Omega = di_X\Omega + i_X d\Omega = 0, \tag{3.1}$$

and the flow at time s

$$\Phi_s\colon \hat{M} \times (-\delta, \delta) \to \hat{M} \times (-\delta, \delta)$$

is of the form

$$\Phi_s(p, t) = (\phi_s(p), t + s).$$

By construction and (3.1), $\phi_s\colon \hat{M} \to \hat{M}$ is a diffeomorphism which satisfies $\phi_s^*\omega_s = \omega_0$. This also shows that ϕ_s is a holomorphic diffeomorphism with respect to the Riemann surface structures M_0 and M_s. □

3.1. Deformations of CMC surfaces.

In the abelian case of tori, it is easy to show that the monodromy representation (based at some fixed point) already determines the conformal type of the immersed CMC torus $f\colon M \to S^3$: The asymptotic behavior of the traces of the monodromies determines the periods of a (non-zero) holomorphic differential on the torus, see [9] for details. Clearly, these periods determine the conformal type of the torus. The main result of this note is the following generalization to compact CMC surfaces of genus $g \geq 2$:

Theorem 3. *Let M be a compact oriented surface. Let $f_t\colon M \to S^3$ be a family of CMC surfaces such that the gauge equivalence classes of associated family of flat connections ${}^t\nabla^\lambda$ are equal for all t and generic $\lambda \in \mathbb{C}^*$. If the CMC surfaces have only simple umbilics (for generic t). Then the induced Riemann surface structures Σ_t are all equivalent and the Hopf differentials coincide.*

Remark 3.2. The theorem remains true for CMC surfaces with periods, i.e., for CMC surfaces into a flat S^3 bundle over M (in the sense of Hitchin's [9] gauge-theoretic harmonic map equations). These are given by solutions (g, Q) of the Gauss-Codazzi equations which are globally defined on the surface M. Here g is a Riemannian metric which induces the spin connection in(2.2), and Q is the Hopf differential. The Gauss-Codazzi equations are then equivalent to the flatness of all connections ∇^λ in (2.2). The theorem also remains true for CMC surfaces (with periods) in the space forms $\mathbb{R}^3$ and $\mathbb{H}^3$.

Proof. We show that there exists closed curves γ_i, $i = 1, ...,$ on $\hat{M}$ which satisfy (2.4) for non-empty sectors and which generate the first homology over $\mathbb{Q}$. Then the proof follows from Lemma 3.1 and Proposition 2.1. We apply Lemma 4.4 in [3]. This Lemma and its proof tell us, that (the pull-back of) Q on $\hat{M}$ is the limit of holomorphic quadratic differentials Q_i with respect to suitable Riemann surface structures $\hat{M}_i$ (note that we identify $\hat{M}_i$ and $\hat{M}$ as smooth surfaces) and which satisfy the following properties:

1. all Q_i have the same number of zeros of the same order as Q;
2. for all $i \in \mathbb{N}$ the horizontal distribution $\mathrm{Re}(\sqrt{Q_i})) = 0$ contains closed curves which do not hit any zeros of Q_i and which generate a Lagrangian subspace $\mathcal{L}$ in the symplectic space $H_1(\hat{M}, \mathbb{R})$;
3. for any Lagrangian subspace $\Lambda \subset H_1(\hat{M}, \mathbb{R})$ the differentials Q_i can be chosen in such a way that $\mathcal{L} \cap \Lambda = \{0\}$;
4. there is an open neighborhood U of the zeros of Q such that $(Q_i)_{|U} = Q_{|U}$ for all $i \in \mathbb{N}$.

We claim that for Q_i close to Q and a closed curve $\gamma \colon S^1 \to \hat{M} \setminus \{\text{zeros of } Q_i\}$ satisfying

$$\mathrm{Re}(\sqrt{Q_i}(\gamma')) = 0$$

condition (2.4) is satisfied for a non-empty sector S. This follows from the fact that $\sqrt{Q_i}_{|U} = \sqrt{Q}_{|U}$ on U and that point-wise on the compact set $\hat{M} \setminus U$ the norms of $\sqrt{Q}$ and $\sqrt{Q_i}$ are bounded from below uniformly. Note that because of (iii) we obtain enough curves γ which generate the first homology $H_1(\hat{M}; \mathbb{Q})$.

Remark 3.3. The statement of the theorem remains true for some classes of CMC surface with higher order umbilics. In particular, the conformal type of a (k, l)-symmetric CMC surface (as defined in [6]) is determined uniquely by the asymptotic behavior (2.5) of the monodromy representation.

Remark 3.4. Combining Theorem 3 with Theorem 7 of [8] shows that deformations of compact CMC surfaces (with simple umbilics) which preserve the conjugacy classes of the monodromy representations of the associated families of flat connections must be induced by a family of dressing transformations. Dressing is a transformation of the family of flat connections induced by a λ-dependent gauge which becomes singular at certain spectral parameters $\lambda_0 \notin S^1 \cup \{0, \infty\}$ where parallel eigenlines with respect to ∇^{λ_0} exist. In the case of higher genus $g \geq 2$, families of dressing transformations can only exists if there is a spectral parameter λ_0 for which ∇^{λ_0} is trivial (up to an diagonal $\mathbb{Z}_2$-gauge). We expect that this would occur only in very exceptional situations.

□

References

[1] A. I. Bobenko, *All constant mean curvature tori in $\mathbb{R}^3$, S^3 ,H^3 in terms of theta-functions*, Math. Ann. 290 (1991), no. 2, 209-245.

[2] M. Fedoryuk, *Asymptotic analysis. Linear ordinary differential equations*, Translated from the Russian by Andrew Rodick. Springer-Verlag, Berlin, 1993.

[3] G. Forni, *Deviation of ergodic averages for area-preserving flows on surfaces of higher genus*, Ann. of Math. (2) 155 (2002), no. 1, 1–103.

[4] D. Gaiotto, G. Moore, A. Neitzke, *Wall-crossing, Hitchin systems, and the WKB approximation*, Adv. Math. 234 (2013), 239–403.

[5] R. Gunning, *Lectures on vector bundles over Riemann surfaces*, University of Tokyo Press, Tokyo; Princeton University Press, Princeton, (1967).

[6] L. Heller, S. Heller, N. Schmitt, *The spectral curve theory for (k,l)-symmetric CMC surfaces*, arxiv: 1503.00969.
[7] S. Heller, *Higher genus minimal surfaces in S^3 and stable bundles,* J. Reine Angew. Math. **685** (2013), 105–122.
[8] S. Heller, *A spectral curve approach to lawson symmetric cmc surfaces of genus* 2*,* Math. Annalen, Volume 360, Issue 3 (2014), Page 607–652.
[9] N. J. Hitchin, *Harmonic maps from a* 2*-torus to the* 3*-sphere*, J. Differential Geom. 31 (1990), no. 3, 627-710.
[10] N. J. Hitchin, *The self-duality equations on a Riemann surface*, Proc. London Math. Soc. (3) 55 (1987), no. 1, 59-126.
[11] A. Katok, B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*, Cambridge Univ. Press, Cambridge, 1995.
[12] M. Kontsevich, A. Zorich, *Connected components of the moduli spaces of Abelian differentials with prescribed singularities*, Invent. Math. 153 (2003), no. 3, 631–678.
[13] W. Wasow, *Asymptotic expansions for ordinary differential equations*, Reprint of the 1976 edition. *Dover Publications, Inc.*, New York, 1987.

Sebastian Heller, Institut für Mathematik, Universität Tübingen, Auf der Morgenstelle 10, 72076 Tübingen, Germany

E-mail address: `heller@mathematik.uni-tuebingen.de`